

SEMI- \mathcal{D} -GROUP COVERINGS OF GROUPS II

JAMES WIEGOLD, Manchester (England)

1. Introduction

The purpose of this note is to confirm two conjectures made in [1]. A group is called * a \mathcal{D} -group if it is the union of non-empty pairwise disjoint proper subsemi-groups, and an \mathcal{S} -group if it has at least one aperiodic homomorphic image. It was proved in [1] that every \mathcal{D} -group is an \mathcal{S} -group, and conjectured that the converse is not true. We shall confirm this by showing that the group G generated by two elements a, b with defining relations

$$a^3 = b^2, \quad ab(a^2b)^5 ab^2 = 1 \tag{1.1}$$

is aperiodic — it was pointed out in [1, page 11] that no group with defining relations like (1.1) can be a \mathcal{D} -group: this fact is very easy to establish. The second derived group $\delta_2(G)$ of G is the cycle generated by b^{16} , so that it is central and in particular G is soluble of length 3.

The other conjecture concerned the existence of a group generated by elements of finite order and having non-periodic centre. Here the construction is considerably easier than that of G : in the group J generated by two elements a, b with defining relations

$$a^3 = b^2 = 1, \quad [[a^2, b], [a, b]], [a] = [[[a^2, b], [a, b]], b] = 1, \tag{1.2}$$

the element $[[a^2, b], [a, b]]$ — obviously central — has infinite order. This element generates the second derived group of J , so that J is also soluble of length 3. I have been unable to construct metabelian examples.

2. The group G

Let us first examine the defining relations

$$a^3 = b^2, \quad aba^2ba^2ba^2ba^2bab^2 = 1$$

* All notation not explained here is to be found in [1].

of G . As a^3 and b^2 are central in G , the second of these can be rewritten as

$$\begin{aligned} baba^2ba^2ba^2bab &= 1, \\ b^{-1}b^2a^{-2}a^3ba^2b^{-1}b^2a^{-1}a^3baa^{-2}a^3b^{-1}b^2a^2ba^{-1}a^3b^{-1}b^2ab &= 1, \\ [b, a^2][b, a][a^2, b][a, b]b^8a^{12} &= 1, \end{aligned}$$

and finally as

$$[a^2, b], [a, b] b^{16} = 1. \quad (2.1)$$

Now the derived group $\delta_1(G)$ of G is generated by all commutators of the form $[a^\lambda, b^\mu]$, where λ and μ are integers; but a^3 and b^2 are central so that λ may be taken modulo 3 and μ modulo 2, which means that $\delta_1(G)$ is generated by $[a^2, b] = u$ and $[a, b] = v$. Relation 2.1 shows that $[u, v]$ is central in G , which gives immediately that $\delta_2(G)$ is central and cyclic, with b^{16} as generator. Further, $\delta_1(G)$ is necessarily second nilpotent — in fact it turns out that is a free second nilpotent group on the generators u, v . We shall show that G is aperiodic by constructing it as a double splitting extension with central amalgamations.

The starting-point is a free second nilpotent group H_0 on two generators, that is, we take a group H_0 by two elements u_0 and v_0 with defining relations

$$[u_0, v_0, u_0] = [u_0, v_0, v_0] = 1. \quad (2.2)$$

The mapping β_0 of the generators of H_0 given by

$$u_0\beta_0 = u_0^{-1}, \quad v_0\beta_0 = v_0^{-1}$$

clearly respects the defining relations (2.2), so it generates an automorphism (also called β_0) of order 2 of H_0 . Let B_0 be the splitting extension* of H_0 by an infinite cycle whose generator b_0 induces β_0 on H_0 :

$$B_0 = GP(u_0, v_0, b_0; [u_0, v_0, u_0] = [u_0, v_0, v_0] = 1, u_0^{b_0} = u_0^{-1}, v_0^{b_0} = v_0^{-1}). \quad (2.3)$$

Then B_0 , as an extension of an aperiodic group by an aperiodic group, is aperiodic; further, $[u_0, v_0]$ and b_0^2 are central, since

$$\begin{aligned} [u_0, v_0]^{b_0} &= [u_0^{-1}, v_0^{-1}] = [u_0, v_0], \\ u_0^{b_0^2} &= (u_0^{-1})^{b_0} = u_0, \quad v_0^{b_0^2} = (v_0^{-1})^{b_0} = v_0. \end{aligned}$$

The first of these relations follows from the first of the identities

$$\begin{aligned} [g^m, h^n] &= [g, h]^{mn} \\ (gh)^m &= g^m h^{m(m-1)/2}, \end{aligned} \quad (2.4)$$

valid for any integers m, n and any elements g, h of a second nilpotent group.

* In general we write $GP(X; R)$ for the group generated by a set X of elements with defining relation R , and $GP(X)$ if R is understood or unimportant.

Clearly, as H_0 intersects the cycle generated by b_0 trivially, the central cycle generated by $[u_0, v_0] b_0^{16}$ intersects H_0 and the cycle $GP(b_0)$ trivially. Thus, if Φ is the canonic homomorphism of B_0 onto $B = B_0/C_0$, then $H = H_0\Phi \simeq H_0$, and $b = b_0\Phi$ has infinite order, so that:

2.5. In the group B generated by three elements u, v, b with defining relations

$$[u, v, u] = [u, v, v] = 1, \quad u^b = u^{-1}, \quad v^b = v^{-1}, \quad [u, v] = b^{-16},$$

the subgroup $H = GP(u, v)$ is free second nilpotent and b has infinite order.

Lemma 2.6. (i) $In B, GP(b) \cap H = GP(b^{16}) = GP([u, v])$.

(ii) B is aperiodic.

Proof. (i) It is clear that $H \cap GP(b) \cong GP(b^{16})$. Let now $x \in H \cap GP(b)$ so that $x = h_0\Phi = b_0^{\lambda}\Phi$ where $h_0 \in H_0$ and λ is an integer. Then $(h_0b_0^{-\lambda})\Phi = 1$ so that $h_0b_0^{-\lambda}$ is a power of $[u_0, v_0] b_0^{16}$, say $[u_0, v_0]^{\epsilon} b_0^{16\epsilon}$; hence $b_0^{-\lambda} = b_0^{16\epsilon}$ so that $x = b^{\lambda} = b^{-16\epsilon} \in GP(b^{16})$. This completes the proof of the first part.

(ii) It is not difficult to see that every element of B can be written in the form

$$g = u^{\alpha}v^{\beta}[u, v]^{\gamma}b^{\delta}$$

with suitable integers $\alpha, \beta, \gamma, \delta$. We distinguish two cases:

(A) If δ is even, b^{δ} is in the centre of B , and, as $[u, v]$ is in the centre of B (both these follow immediately from the fact that b_0^2 is central in B_0), then for any integer n ,

$$g^n = (u^{\alpha}v^{\beta})^n [u, v]^{\gamma n} b^{n\delta}$$

But $(u^{\alpha}v^{\beta})^n = u^{\alpha n}v^{\beta n}[u, v]^{\epsilon}$ for some ϵ , by 2.4; hence

$$g^n = u^{\alpha n}v^{\beta n}[u, v]^{\epsilon} b^{n\delta}$$

for some ϵ . If $g^n = 1$ for some $n \neq 0$, then, as $[u, v] = b^{-16}$, $u^{\alpha n}v^{\beta n}$ lies in the cycle generated by b . The first part of the lemma then gives that $u^{\alpha n}v^{\beta n}$ lies in the cycle generated by $[u, v]$, so that since H is free second nilpotent on u and v , $n\alpha = n\beta = 0$. Consequently $\alpha = \beta = 0$, $g = [u, v]^{\gamma}b^{\delta}$, so g lies in the cycle generated by b ; and this means it can have finite order if and only if it is the unit element.

(B) If δ is odd, say $\delta = 2\delta' + 1$, then

$$g = u^{\alpha}v^{\beta}b[u, v]^{\gamma}b^{2\delta'}.$$

so that

$$\begin{aligned} g^2 &= u^{\alpha}v^{\beta}bu^{\alpha}v^{\beta}b[u, v]^2b^{4\delta'} \\ &= u^{\alpha}v^{\beta}(u^{\alpha}v^{\beta})^{b^{-1}}b^2[u, v]^2b^{4\delta'} \\ &= u^{\alpha}v^{\beta}u^{-\alpha}v^{-\beta}[u, v]^2b^{4\delta'+2}, \end{aligned}$$

using throughout the fact that b^2 is central. Finally

$$\begin{aligned} g^2 &= [u^{-\alpha}, v^{-\beta}] [u, v]^{2\gamma} b^{4\delta+2} \\ &= [u, v]^{2\gamma+\alpha\beta} b^{4\delta+2} \\ &= b^{-1(6(2\gamma+\alpha\beta)+4\delta+2)} \\ &= b^{4\delta'+2-16\epsilon}. \end{aligned} \quad (\text{by 2.4})$$

For some integer ϵ . Now $4\delta' + 2 - 16\epsilon \neq 0$ since $2\delta' + 1 \neq 8\epsilon$; hence g^2 has infinite order and some must g .

Thus we have proved that only the unit element of B is of finite order, and B is aperiodic.

It will turn out that B is the normal closure of b in the original group G (we shall not prove this) — hence the use of the letter b in defining B .

The next stage is to form a splitting extension of B by an infinite cycle. For clarity at this point we take an isomorphic copy B_1 of B with generators u_1, v_1, b_1 and defining relations analogous to those given at 2.5. It is a matter of simple routine to verify that the mapping α_1 of the generators of B_1 defined by

$$u_1\alpha_1 = v_1^{-1}, \quad v_1\alpha_1 = u_1v_1^{-1}, \quad b_1\alpha_1 = b_1v_1^{-1}$$

respects the defining relations of B_1 . Thus α_1 generates an endomorphism (also called α_1) of B_1 . Further, α_1^3 is the identity automorphism, as

$$\left. \begin{aligned} u_1\alpha_1^2 &= (v_1\alpha_1)^{-1} = v_1u_1^{-1}, & u_1\alpha_1^3 &= v_1\alpha_1(u_1\alpha_1)^{-1} = u_1v_1^{-1}v_1 = u_1; \\ v_1\alpha_1^2 &= (u_1v_1^{-1})\alpha_1 = v_1^{-1}v_1u_1^{-1} = u_1^{-1}, & v_1\alpha_1^3 &= (u_1\alpha_1)^{-1} = v_1; \\ b_1\alpha_1^2 &= b_1\alpha_1(v_1\alpha_1)^{-1} = b_1v_1^{-1}v_1u_1^{-1} = b_1u_1^{-1}, \\ b_1\alpha_1^3 &= b_1\alpha_1(u_1\alpha_1)^{-1} = b_1v_1^{-1}v_1 = b_1. \end{aligned} \right\} \quad (2.7)$$

This means that α_1 has a two-sided inverse and is consequently an automorphism of order 3 of B_1 . Form the splitting extension of B by an infinite cycle whose generator a_1 induces α_1 on B_1 : this is a group G_1 generated by four elements u_1, v_1, b_1, a_1 with defining relations

$$\left. \begin{aligned} [u_1, v_1, u_1] &= [u_1, v_1, v_1] = 1, & u_1^{b_1} &= u_1^{-1}, & v_1^{b_1} &= v_1^{-1}, \\ b_1^{-16} &= [u_1, v_1, \underbrace{a_1^{-1}}_{\text{central}}] = v_1^{-1}, & v_1^{a_1} &= u_1v_1^{-1}, & b_1^{a_1} &= b_1v_1^{-1}. \end{aligned} \right\} \quad (2.8)$$

Again G_1 is aperiodic and it is routine to verify that b_1^2 and a_1^3 are central in G_1 . Let A_1, C_1 be the cycles generated by a_1 and $b_1^{-2}a_1^3$ respectively. Then C_1 is central in G_1 and misses B_1 and A_1 so that if ψ is the canonic homomorphism of G_1 onto $G_0 = G_1/C_1$, then $B_1\psi \simeq B_1$ and $a_1\psi = a$ has infinite order. Thus we have proved the first part of the following lemma — the second is proved in a manner completely analogous to that of Lemma 2.6 (i):

Lemma 2.9. *In the group G_0 generated by elements u, v, b, a with defining relations*

$$\begin{aligned} [u, v, u] &= [u, v, v] = 1, & u^b &= u^{-1}, & v^b &= v^{-1}, & b^{-16} &= [u, v], \\ u^a &= v^{-1}, & v^a &= uv^{-1}, & b^a &= bu^{-1}, & a^3 &= b^2, \end{aligned}$$

the subgroup $B = Gp(u, v, b)$ is aperiodic and $A = Gp(a)$ is infinite.

(ii) $B \cap A = Gp(a^3) = Gp(b^2)$.

We can now prove the main result.

Theorem 2.10. G_0 is aperiodic.

Proof. Since B is normal in G_0 and $a^3 \in B$, every element of G_0 can be put in one of the forms h, ha, ha^2 with $h \in B$. Obviously, if h is of finite order, $h = 1$ because B is aperiodic. We shall show first that every element of the form ha is of infinite order. As G/B is of order 3 so that $(ha)^3 \in B$, the element ha can have finite order if and only if $(ha)^3 = 1$. Now

$$\begin{aligned} (ha)^2 &= hah a = hah a^{-1} a^2 = hh a^{-1} a^2, \\ (ha)^3 &= hh a^{-1} a^2 ha = hh a^{-1} a^2 ha^{-2} a^3 \\ &= hh a^{-1} h a^{-2} a^3 \\ &= hh a^{-1} h a^{-2} b^2. \end{aligned}$$

It is easy to check that h can be expressed in the form

$$h = u^x v^y [u, v]^z b^d$$

for suitable integers $\alpha, \beta, \gamma, \delta$. Once again there are two cases, depending on the parity of δ .

(A) If δ is even, b^d is central so that as a^3 is central,

$$\begin{aligned} hh a^{-1} h a^{-2} b^2 &= hh a^3 h a^2 b^2 \\ &= u^{2x} v^{2y} (u^x v^y a^3)^2 (u^x v^y)^2 [u, v]^{3z} b^{3d+2} \\ &= u^{2x} v^{2y} (uv^{-1})^x u^{-\beta} v^{-\alpha} (uv^{-1})^y [u, v]^{3z} b^{3d+2} \\ &= u^{2x} v^{2y} u^{-\alpha} u^{-\beta} v^{-\alpha} u^{-\beta} v^{-\beta} [u, v]^{3z+\Phi(\delta)-\Phi(\alpha)} b^{3d+2}, \end{aligned} \quad (\text{from 2.7})$$

this from 2.4, where for any integer $k, \Phi(k) = k(k-1)/2$. A little computation now gives

$$hh a^{-1} h a^{-2} b^2 = [u, v]^{3z+\Phi(\delta)-\Phi(\alpha)+2\alpha\beta} b^{3d+2}.$$

If this is 1, then as $[u, v] = b^{-16}$ and b has infinite order,

$$3\delta + 2 = 16(3\gamma + \Phi(\beta) - \Phi(\alpha) + \alpha^2 + 2\alpha\beta).$$

But $\delta = 2\delta'$ so that

$$3\delta' + 1 = 8(3\gamma + \Phi(\beta) - \Phi(\alpha) + \alpha^2 + 2\alpha\beta).$$

We shall obtain a contradiction by showing that the right hand side of this equation is never congruent to 1 modulo 3. First,

$$\begin{aligned} 8(3\gamma + \Phi(\beta) - \Phi(\alpha) + \alpha^2 + 2\alpha\beta) &\equiv 2(\Phi(\beta) - \Phi(\alpha) + \alpha^2 + 2\alpha\beta) \pmod{3} \\ &\equiv \beta(\beta - 1) - \alpha(\alpha - 1) + 2\alpha^2 + \alpha\beta \pmod{3} \\ &= \beta^2 + \alpha^2 + \alpha - \beta + \alpha\beta. \end{aligned}$$

The rest is routine: one verifies that $\beta^2 + \alpha^2 + \alpha - \beta + \alpha\beta$ is always congruent to 0 or 2 modulo 3, in all the 9 cases arising out of the 3 congruences modulo 3 possible for each of α and β .

Hence if δ is even, ha is of infinite order.

(B) If δ is odd, say $\delta = 2\delta' + 1$, then

$$h = u^{\delta'} v^{\delta'} b z,$$

where $z = [u, v] b^{2\delta'}$ is in the centre of G_0 . Then

$$h h^{\alpha^2} h^{\beta^2} = u^{\delta'} v^{\delta'} b (u^{\alpha^2})^{\delta'} (v^{\alpha^2})^{\delta'} b^{\delta'} (u^{\beta^2})^{\delta'} (v^{\beta^2})^{\delta'} b^{2\delta'},$$

and consideration of the image of this element in $G_0/GP(u, v)$ shows it (the original element) to be of the form $v^{\delta'} b^{5+6\delta'}$ for some $v' \in GP(u, v)$. If it is 1, then by lemma 2.6(i), $b^{5+6\delta'}$ lies in $GP(b^{16})$. This means that $5 + 6\delta'$ is divisible by 16, which is a manifest contradiction. Thus $h h^{\alpha^2} h^{\beta^2}$ is never the unit element, and it follows that ha has infinite order.

The proof concludes with the remark that ha^2 must also have infinite order, since

$$\begin{aligned} (ha^2)^2 &= ha^2 ha^2 = h h^{\alpha^2} a^4 \\ &= h h^{\alpha^2} a^3 a \end{aligned}$$

is of the form $h'a$ with $h' \in B$.

It remains only to show that G_0 is isomorphic with G . The defining relations of G_0 in terms of the generators u, v, a, b are

$$\begin{aligned} [u, v, u] &= [u, v, v] = 1, & u^b &= u^{-1}, & v^b &= v^{-1}, \\ u^a &= v^{-1}, & v^a &= uv^{-1}, & b^a &= bv^{-1}, \\ b^{-16} &= [u, v], & a^3 &= b^2. \end{aligned}$$

From these it follows straight away that

$$\begin{aligned} v &= (b^{-1})^{16} = [a, b] \\ u &= v^a = [a, b]^a [a, b] = [a^2, b] \end{aligned}$$

so that G_0 is generated by a and b . One now readily verifies that the first 8 defining relations of G_0 are consequences of the last two; these are precisely the relations 2.1, so that in fact G and G_0 are isomorphic.

To sum up, define a \mathbb{C} -group to be one whose second derived group is central. With the notation of [1],

Theorem 2.11 $[R] \cap [C] \supseteq [D] \cap [G]$.

3. The group J

Here we shall only state results: the proofs are a matter of simple routine. We again start with a free second nilpotent group on two generators:

$$H = GP(u, v; [u, v, u] = [u, v, v] = 1),$$

and form the splitting extension of H by a cycle of order 2 whose generator b induces the automorphism of order 2 of H generated by the mappings $u \rightarrow u^{-1}, v \rightarrow v^{-1}$;

$$H = GP(u, v, b; [u, v, u] = 1, b^2 = 1, u^b = u^{-1}, v^b = v^{-1}).$$

In this u and v still generate a free second nilpotent group, so in particular $[u, v]$ is of infinite order. Next form the splitting extension J of B be a cycle of order 3 whose generator a induces the automorphism of B defined by the mappings

$$u \rightarrow v^{-1}, \quad v \rightarrow uv^{-1}, \quad b \rightarrow bv^{-1};$$

J is generated by u, v, a, b with the defining relations of B together with

$$a^3 = 1, \quad u^a = v^{-1}, \quad v^a = uv^{-1}, \quad b^a = bv^{-1}.$$

Note (as with G) that $v = [a, b]$, $u = [a^2, b]$ so that J is generated by a and b , and that $[u, v]$ is central, as

$$\begin{aligned} [u, v]^a &= [u^{-1}, v^{-1}] = [u, v] \\ [u, v]^b &= [v^{-1}, uv^{-1}] = [v^{-1}, u] = [u, v]. \end{aligned}$$

This completes the example, except for the simple verification that the group J constructed here is in fact that mentioned in the introduction.

REFERENCES

[1] Wieland James, Semigroup coverings of groups, *Mathematicko-fizikalny časopis SAV* 11 (1961), 3-13.

Received January 27, 1962.

*Department of Mathematics,
Manchester College of Science and Technology,
Manchester 1, England*

ПОКРЫТИЕ ГРУПП ПОЛУГРУППАМИ

Джеймс Уайланд

Резюме

Настоящая статья является продолжением статьи (1). В статье (1) введено понятие \mathbb{D} -группы (это группа, являющаяся объединением попарно непересекающихся собственных подполугрупп) и понятие \mathbb{S} -группы (это группа, которая имеет хотя бы один алгебраический гомоморфный образ) и доказывается, что всякая \mathbb{D} -группа является \mathbb{S} -группой. В настоящей статье построена группа G , которая является \mathbb{S} -группой, но не является \mathbb{D} -группой. Кроме того, построен пример группы с образующими конечного порядка, центр которой не является периодической группой.