

FRACTIONAL SUPERSYMMETRY THROUGH GENERALIZED ANYONIC ALGEBRA**J. Douari^{ab1}, Y. Hassouni^{b2}***(a) The Abdus Salam ICTP - Strada Costiera 11, 34100 Trieste, Italy**(b) Faculté des Sciences, Département de Physique, LPT-ICAC
Av. Ibn Battouta, B.P.1014, Agdal, Rabat, Morocco*

Received 4 April 2000, in final form 22 March 2001, accepted 4 April 2001

The construction of anyonic operators and algebra is generalized by using quons operators. Therefore, the particular version of fractional supersymmetry is constructed on the two-dimensional lattice by associating two generalized anyons of different kinds. The fractional supersymmetry Hamiltonian operator is obtained on the two-dimensional lattice and the quantum algebra $U_q(sl_2)$ is realized.

PACS: 05.30.-d, 74.20.-z, 74.65.+n

1 Introduction

A growing interest has been recently devoted to fractional statistics [1, 2] that describe particles called anyons, with fractional spin [3–5]. These anyons interpolate between bosons and fermions. Furthermore that have been shown to be non-local particles defined on the two-dimensional space. Mathematically, the group symmetry associated to the anyonic systems involves some special Lie algebras appearing as a quantum deformation of the usual Lie algebras and Lie groups [6–10]. Indeed, one can prove that quantum groups [11] are the mathematical objects allowing the description of these particular systems [12, 13]. Many works have been devoted, for the last few years, to the construction and the realization of quantum groups. We note, for example, that the anyonic algebra [14] leads to the obtaining of these quantum algebras. The anyonic operators are introduced on the two-dimensional lattice as non-local operators seen as a generalization of generators of the Jordan-Wigner transformation [15].

The present work is concerned the study of N=2 fractional supersymmetry (FSUSY) [16–19] on the two-dimensional lattice. We obtain exactly the same algebraic structure of latter starting from an introduction of the generalized anyonic algebra. Then they introduce some special operators which can be seen as generalized anyonic operators. We notice that this construction is different from the one used to reproduce the N=2 FSUSY algebra based on the quonic algebra [19]. This algebraic study allows us to find the Hamiltonian operator describing one N=2 FSUSY system.

¹E-mail address: douarij@ictp.trieste.it²E-mail address: y-hassou@fsr.ac.ma

This paper is organized as follows : In section 2, as a generalization, we construct new exotic operators by using quonic operators as basic ones. Therefore we realize the generalized anyonic operators and the corresponding algebra. In the third section, the last new operators and algebra are used to construct N=2 FSUSY on the two-dimensional lattice corresponding to anyonic systems as a certain coupling of two generalized anyonic oscillators of different kinds (γ and δ) [14]. We proceed as in the work of [19] where we have constructed the N=2 FSUSY through two different quons. Another result consists of the realization of the quantum algebra $U_q(sl_2)$ using the supercharges already constructed. Finally, section 4 presents some concluding remarks.

2 Generalized anyonic algebra

As a reminder, the anyonic operators are seen as non-local two-dimensional operators interpolating between bosonic and fermionic ones. The generalization of these operators allows us to introduce the generalized anyons. We will use the famous angle function $\Theta(x, y)$ appearing in the construction and the description of anyons in the work of [6].




Fig.1

B

First, here is a brief review on this angle function. One designs by γ_x the cut associated to each point x that we denote by x_γ on the two-dimensional lattice Ω . Denoting by Ω^* the dual lattice of Ω ; it is a set of points $x^* = x + o^*$ with $o^* = (\frac{1}{2}\epsilon, \frac{1}{2}\epsilon)$ the origin of Ω^* and ϵ its spacing which eventually will be sent to zero. In this case γ_x will be on Ω^* from minus infinity to x^* along the x -axis (Fig.1). One considers another type of cut. We choose the set of cuts δ_x coming from plus infinity to $^*x = x - o^*$ along the x -axis (Fig.2). Consequently, the two types of cuts γ_x and δ_x involve an ordering and opposite ordering respectively of points x on Ω . This is described by the following proposition

$$x_\delta < y_\delta \Leftrightarrow x_\gamma > y_\gamma \Leftrightarrow x > y \Leftrightarrow \begin{cases} x_2 > y_2, \\ x_1 > y_1, x_2 = y_2 \end{cases} \tag{1}$$




Fig.2

 \tilde{B}

Owing to equation (1) the angle functions satisfy

$$\begin{aligned} \Theta_{\gamma_x}(x, y) - \Theta_{\gamma_y}(y, x) &= \begin{cases} \pi, & x > y \\ -\pi, & x < y \end{cases} \\ \Theta_{\delta_x}(x, y) - \Theta_{\delta_y}(y, x) &= \begin{cases} -\pi, & x > y \\ \pi, & x < y \end{cases} \\ \Theta_{\delta_x}(x, y) - \Theta_{\gamma_x}(x, y) &= \begin{cases} -\pi, & x > y \\ \pi, & x < y \end{cases} \\ \Theta_{\delta_x}(x, y) - \Theta_{\gamma_y}(y, x) &= 0, \quad \forall x, y \in \Omega. \end{aligned} \quad (2)$$

Now let us introduce the operators $K_i(x_\alpha)$ those called the disorder ones. They are expressed by

$$K_i(x_\alpha) = \exp \left\{ i\nu \sum_{y \neq x} \Theta_{\alpha_x}(x, y) \left[N_i(y) - \frac{1}{2} \right] \right\} \quad (3)$$

with $\alpha_x = \gamma_x$ or δ_x and $i = 1, 2, \dots, n, n \in \mathbf{N}$. In the equality (3) $N_i(y)$ is nothing but the number operator of quons on the two-dimensional lattice, defined by the generators $a_i^\dagger(x)$ and $a_i(x)$ as follows

$$\begin{aligned} a_i^\dagger(x) a_i(x) &= [N_i(x)]_{q_i} \\ a_i(x) a_i^\dagger(x) &= [N_i(x) + 1]_{q_i} \end{aligned} \quad (4)$$

where $[x]_q = \frac{q^x - 1}{q - 1}$. The operators $a_i^\dagger(x)$ and $a_i(x)$ are respectively the creation and annihilation quonic operators on the two-dimensional lattice Ω , satisfying the relations

$$\begin{aligned} [a_i(x), a_j^\dagger(y)]_{q_i} &= \delta_{ij} \delta(x, y) \\ [a_i(x), a_j(y)]_{q_i} &= 0 \quad \forall x, y, \forall i, j \\ [a_i^\dagger(x), a_j^\dagger(y)]_{q_i} &= 0 \quad \forall x, y, \forall i, j \\ [N_i(x), a_j(y)] &= -\delta_{ij} \delta(x, y) a_i(x) \\ [N_i(x), a_j^\dagger(y)] &= \delta_{ij} \delta(x, y) a_i^\dagger(x) \end{aligned} \quad (5)$$

The lattice Dirac function is defined by

$$\delta(x, y) = \begin{cases} 1 & \text{if } x = y \\ 0 & \text{if } x \neq 0 \end{cases} \quad (6)$$

In the relations (5), one can consider the operators $a_i^\dagger(x)$, $a_i(x)$ and $N_i(x)$; those are generators of the oscillator algebra describing a system of quons. They are seen as non-local particles and we define them to satisfy the equality

$$(a_i(x))^{d_i} = (a_i^\dagger(x))^{d_i} = 0 \quad (7)$$

where we take the deformation parameter q_i to be a root of unity, $q_i^{d_i} = 1$ then $q_i = e^{i\frac{2\pi}{d_i}}$. One can show that the irreducible representation space (Fock space) of the algebra (relations (5)) is given by the set

$$F_{i_x} = \{|n_{i_x}\rangle, n_{i_x} = 0, 1, \dots, d_i - 1\} \quad (8)$$

where the notation i_x means that this Fock space is introduced in each site of the Ω . The actions of the generators $a_i(x)$, $a_i^\dagger(x)$ and $N_i(x)$ are expressed by the following equalities

$$\begin{aligned} a_i^\dagger(x)|n_{i_x}\rangle &= |n_{i_x} + 1\rangle, & a_i^\dagger(x)|d_i - 1\rangle &= 0 \\ a_i(x)|n_{i_x}\rangle &= [n_{i_x}]_{q_i}|n_{i_x} - 1\rangle, & a_i(x)|0\rangle &= 0. \end{aligned} \quad (9)$$

Now we introduce an operator allowing the description of the N=2 FSUSY on the two-dimensional lattice. This realization is seen as an anyonic one, in the sense that the generators will depend on a function which seems to be the angle function used by the authors in the work of [14] when they describe the system of anyons. This leads to the definition of the operators

$$A_i(x_\alpha) = K_i(x_\alpha)a_i(x), \quad (10)$$

where $K_i(x_\alpha)$ is the disorder operator introduced in equation (3).

One proves that these operators obey the following commutation relations

$$\begin{aligned} [A_i(x_\gamma), A_i(y_\gamma)]_{q_i p^{-1}} &= 0, \quad x > y \\ [A_i^\dagger(x_\gamma), A_i^\dagger(y_\gamma)]_{q_i p^{-1}} &= 0, \quad x > y \\ [A_i(x_\gamma), A_i^\dagger(y_\gamma)]_{q_i p} &= 0, \quad x > y \\ [A_i^\dagger(x_\gamma), A_i(y_\gamma)]_{q_i p} &= 0, \quad x > y \\ [A_i(x_\gamma), A_i^\dagger(x_\gamma)]_{q_i} &= 1 \\ [A_i(x_\gamma), A_j(y_\gamma)] &= 0, \quad \forall i \neq j, \forall x, y \in \Omega \\ [A_i^\dagger(x_\gamma), A_j^\dagger(y_\gamma)] &= 0, \quad \forall i \neq j, \forall x, y \in \Omega \\ [A_i^\dagger(x_\gamma), A_j(y_\gamma)] &= 0, \quad \forall i \neq j, \forall x, y \in \Omega \\ [A_i(x_\gamma), A_j^\dagger(y_\gamma)] &= 0, \quad \forall i \neq j, \forall x, y \in \Omega. \end{aligned} \quad (11)$$

for the anyonic operators of type γ , where $p = e^{i\nu\pi}$. We point out that the same results can be obtained for the kind δ in replacing p by p^{-1} in (11). It is obvious thus to get the commutation relations between the different kinds of generalized anyonic operators, so we have

$$\begin{aligned} [A_i(x_\delta), A_j(y_\gamma)]_{q_i}^{\delta_{ij}} &= 0, \quad \forall x, y \in \Omega \\ [A_i(x_\delta), A_j^\dagger(y_\gamma)]_{q_i}^{\delta_{ij}} &= \delta_{ij} \delta(x, y) p^{-\left(\sum_{z < x} - \sum_{z > x}\right) [N_i(z) - \frac{1}{2}]} \end{aligned} \quad (12)$$

and ν is seen as the statistical parameter [1, 2].

By considering the generalized anyonic oscillators $A_i(x_\alpha)$ and $A_i^\dagger(x_\alpha)$, we can demonstrate the following relations

$$(A_i(x_\alpha))^{d_i} = (A_i^\dagger(x_\alpha))^{d_i} = 0 \quad (13)$$

with $\alpha = \gamma, \delta$ and $i = 1, 2, \dots, N$. Equation (13) can be seen as a generalization of the hard-core condition found when we studied the generalized statistics in the works [20–22]. Indeed, for $d_i = 2$ one recovers the result allowing us, in this work, to think about some connection between the generalized statistics and the anyonic ones. Returning to the present paper, the relation (13) can be seen a nilpotency condition leading to the study of quons on the two-dimensional lattice. This equality is obtained starting from the function $K_i(x_\alpha)$ discussed in the work of [14], which satisfies the following equations

$$\begin{aligned} K_i^\dagger(x_\alpha) a_i(y) &= \exp[i\nu\Theta_{\alpha_x}(x, y)] a_i(y) K_i^\dagger(x_\alpha) \\ K_i^\dagger(x_\alpha) a_i^\dagger(y) &= \exp[-i\nu\Theta_{\alpha_x}(x, y)] a_i^\dagger(y) K_i^\dagger(x_\alpha) \\ K_i(x_\alpha) a_i(y) &= \exp[-i\nu\Theta_{\alpha_x}(x, y)] a_i(y) K_i(x_\alpha) \\ K_i(x_\alpha) a_i^\dagger(y) &= \exp[i\nu\Theta_{\alpha_x}(x, y)] a_i^\dagger(y) K_i(x_\alpha) \\ K_i^\dagger(x_\alpha) K_i(y_\alpha) &= K_i(y_\alpha) K_i^\dagger(x_\alpha) \end{aligned} \quad (14)$$

It can be proved that, on the above Fock space, these algebraic relations are consistent with the following equalities

$$\begin{aligned} A_i^\dagger(x_\alpha) |n_{i_x} \rangle &= \exp\left[i\frac{\nu}{2} \sum_{y \neq x} \Theta_{\alpha_x}(x, y)\right] |n_{i_x} + 1 \rangle \\ A_i(x_\alpha) |n_{i_x} \rangle &= [n_{i_x}]_{q_i} \exp\left[-i\frac{\nu}{2} \sum_{y \neq x} \Theta_{\alpha_x}(x, y)\right] |n_{i_x} - 1 \rangle \\ A_i^\dagger(x_\alpha) |d_i - 1 \rangle &= 0 \\ A_i(x_\alpha) |0 \rangle &= 0. \end{aligned} \quad (15)$$

We note that the operators $A_i^\dagger(x_\alpha)$ and $A_i(x_\alpha)$ are seen, owing to relations (14) and (15), as generalized creation and annihilation anyonic operators respectively.

3 N=2 FSUSY on the two-dimensional lattice

The construction of generalized anyonic operators allows us to realize the N=2 FSUSY. This realization involves two different generalized anyons and is analogous to the one obtained via two different quons in the work of [19]. We introduce then the fractional supercharges on Ω as

$$\begin{aligned} Q_+(x) &= A_1^\dagger(x_\gamma) A_2(x_\delta) \\ Q_-(x) &= A_1(x_\delta) A_2^\dagger(x_\gamma). \end{aligned} \quad (16)$$

basing on the equality (13); one can get

$$(Q_\pm(x))^{d_2} = 0, \quad (17)$$

where d_2 is required to be $< d_1$.

In using the construction given in the relation (16), we can get the action of the above fractional supercharges on the Fock space defined by the tensor product

$$F_x = F_{1_x} \otimes F_{2_x} \quad (18)$$

with F_{1_x} and F_{2_x} corresponding respectively to the different generalized anyons used to introduce the fractional supercharges.

Now we can easily get the relations

$$\begin{aligned} Q_+(x)|n_{1_x} \rangle \otimes |n_{2_x} \rangle &= p^{\frac{1}{2}} [n_{2_x}]_{q_2} |n_{1_x} + 1 \rangle \otimes |n_{2_x} - 1 \rangle \\ Q_-(x)|n_{1_x} \rangle \otimes |n_{2_x} \rangle &= p^{\frac{1}{2}} [n_{1_x}]_{q_1} |n_{1_x} - 1 \rangle \otimes |n_{2_x} + 1 \rangle \end{aligned} \quad (19)$$

Through this realization of the fractional supercharges $Q_{\pm}(x)$, we can show that these new operators satisfy the following commutation relation in the case of $x > y$ on Ω

$$q_1 Q_+(x) Q_-(y) - q_2 Q_-(y) Q_+(x) = \delta(x, y) P [q_1 [N_1(x)]_{q_1} - q_2 [N_2(x)]_{q_2}] \quad (20)$$

where the operator P is written as

$$P = p \left(\sum_{z < x} - \sum_{z > x} \right) [N_1(z) + N_2(z) - 1] \quad (21)$$

One can remark that the equality (20) is not invariant under the hermitian conjugation. This is related to the fact that these generators involve complex numbers $q_{1,2}$ and p which are different from ± 1 . To avoid this difficulty we introduce the hermitian conjugate of the generators $Q_{\pm}(x)$ and after calculation it is easy to obtain the conjugate equation of (20) as

$$\begin{aligned} q_1^{-1} Q_-^\dagger(y) Q_+^\dagger(x) - q_2^{-1} Q_+^\dagger(x) Q_-^\dagger(y) &= \delta(x, y) P^{-1} \\ &\times [q_1^{-1} [N_1(x)]_{q_1^{-1}} - q_2^{-1} [N_2(x)]_{q_2^{-1}}], \end{aligned} \quad (22)$$

We can also verify that

$$(Q_{\pm}^\dagger(x))^{d_2} = 0. \quad (23)$$

for $d_2 < d_1$.

Another reason to introduce this operation is to construct the Hamiltonian operator corresponding to this system. We can thus express the FSUSY Hamiltonian operator as

$$\begin{aligned} q_1 Q_+(x) Q_-(y) + q_1^{-1} Q_-^\dagger(y) Q_+^\dagger(x) - q_2 Q_-(y) Q_+(x) - q_2^{-1} Q_+^\dagger(x) Q_-^\dagger(y) \\ = \delta(x, y) [P q_1 [N_1(x)]_{q_1} + P^{-1} q_1^{-1} [N_1(x)]_{q_1^{-1}} \\ - P q_2 [N_2(x)]_{q_2} - P^{-1} q_2^{-1} [N_2(x)]_{q_2^{-1}}]. \end{aligned} \quad (24)$$

Using the relation

$$[N_i(x)]_{q_i^{-1}} = q_i^{1-N_i(x)} [N_i(x)]_{q_i}, \quad (25)$$

the RHS of (24) can be rewritten as

$$RHS = \delta(x, y) [P^{-1} q_1^{-N_1(x)} + P q_1] [N_1(x)]_{q_1} - [P^{-1} q_2^{-N_2(x)} + P q_2] [N_2(x)]_{q_2}, \quad (26)$$

The last equation can be interpreted as a Hamiltonian of the generalized anyonic system investigated. So, we can rewrite this Hamiltonian operator as

$$H(x) = \sum_{i,j=1,2} \frac{\epsilon_{ij}}{\sin(\pi/d_i)} \left\{ \sin \left[\nu\pi \sum_{z \in \Omega} \aleph(z) + \frac{\pi}{d_i} [2N_i(x) + 1] \right] - \sin \left[\nu\pi \sum_{z \in \Omega} \aleph(z) + \frac{\pi}{d_i} \right] \right\}, \quad (27)$$

where $(\epsilon_{ij}) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ and $\sum_{z \in \Omega} \aleph(z) = (\sum_{z < x} - \sum_{z > x})(N_1(z) + N_2(z) - 1)$. Up to now, we can recapulate our result which consists on the complete description of the N=2 FSUSY basing on a given anyonic realization.

Furthermore, a global version of this realization can be readily constructed as follows

$$H = \sum_{x \in \Omega} H(x), \quad (28)$$

where the global supercharges are defined as

$$Q_{\pm} = \sum_{x \in \Omega} Q_{\pm}(x), \quad (29)$$

In addition, we can link the N=2 FSUSY obtained on the two-dimensional lattice to the "local" algebra $U_q(sl_2)$ in considering $q_1 = q_2 = q$. In this particular case we define the three local generators as

$$\begin{aligned} J_{\pm}(x) &= P^{-\frac{1}{2}} q^{-\frac{N_2(x)}{2}} Q_{\pm}(x) \\ J_3(x) &= \frac{1}{2}(N_1(x) - N_2(x)). \end{aligned} \quad (30)$$

We can easily check that these local densities of quantum group generators satisfy the following commutation relations

$$\begin{aligned} [J_+(x), J_-(y)] &= \delta(x, y) [2J_3(x)]_q \\ [J_3(x), J_{\pm}(y)] &= \pm \delta(x, y) J_{\pm}(x). \end{aligned} \quad (31)$$

Thus to define the global generators it is sufficient to write

$$\begin{aligned} J_{\pm} &= \sum_{x \in \Omega} J_{\pm}(x) \\ J_3 &= \sum_{x \in \Omega} J_3(x) \end{aligned} \quad (32)$$

and close the $U_q(sl_2)$ algebra as

$$\begin{aligned} [J_+, J_-] &= [2J_3]_q \\ [J_3, J_{\pm}] &= \pm J_{\pm}. \end{aligned} \quad (33)$$

Consequently, we have closed the algebra of $U_q(sl_2)$ generated by J_{\pm} and J_3 which are built out of generalized anyonic oscillators.

4 Conclusion

To conclude, we can summarize the lines of this paper in saying that we have constructed generalized anyonic operators on the two-dimensional lattice by using the q -bosonic operators or called the so-quonic operators, as a generalization of ones defined in the paper [5]. Moreover, we have realized the $N=2$ FSUSY on the two-dimensional lattice, where the supercharges are constructed by coupling two different generalized anyonic operators, and the FSUSY Hamiltonian operator of the corresponding system is given. Thus from the $N=2$ FSUSY realized the well-known algebra $U_q(sl_2)$ is derived.

Acknowledgments The author J. Douari would like to thank the Max-Planck-Institut für Physik Komplexer Systeme for link hospitality during the stage in which one part of this paper was done and the International Atomic Energy Agency and UNESCO for hospitality at the Abdus Salam International Centre for Theoretical Physics, Trieste. This work was achieved within the framework of the Associateship Scheme of the Abdus Salam International Centre for Theoretical Physics. And special thanks go to Professors Manfred Scheunert, Ruibin Zhang and G. Thompson for inspiring comments and many useful discussions.

References

- [1] J. F. Schonfeld: *Nucl. Phys. B* **185** (1981) 127
- [2] F. Wilczek: *Phys. Rev. Lett.* **48** (1982) 1144 ; R. B. Laughlin: *Science* **242** (1988) 525
- [3] J. M. Leinaas, J. Myrheim: *Nuovo Cimento B* **37** (1977) 1 ; J. M. Leinaas: *Symmetry and Structural Properties of Condensed Matter*, Poznan 1992. World Sci., Singapore, 1993
- [4] A. Lerda: *Anyons: Quantum Mechanics of Particles with Fractional Statistics*, Springer-Verlag, Berlin, Germany 1992
- [5] I. J. R. Aitchison, N. E. Mavromatos: *Contemp. Phys.* **32** (1991) 219
- [6] T. Brzezinski, I. L. Egusquiza, A. J. Macfarlane: *Phys. Lett. B* **311** (1993) 202
- [7] A. J. Macfarlane: *J. Phys. A* **22** (1989) 4581
- [8] M. Chaichan: *Phys. Lett. B* **234** (1990) 72
- [9] L. C. Biedenharn: *J. Phys. A* **22** (1989) L873
- [10] M. A. Vasiliev: *Int. J. Mod. Phys. A* **6** (1991) 1115
- [11] A. P. Isaev: *Fiz. El'em. Chastits At. Yadra* **26** (1995) 1204
- [12] M. Frau, M. A. R. Monteiro, S. Sciuto: *J. Phys. A: Math. Gen.* **27** (1994) 801
- [13] L. Frappat, A. Sciarrino, S. Sciuto, P. Sorba: *Phys. Lett. B* **369** (1996) 313
- [14] A. Lerda, S. Sciuto: *Nucl. Phys. B* **401** (1993) 613
- [15] P. Jordan, E. P. Wigner: *Z. Phys. B* **47** (1928) 631
- [16] A. Lerda, S. Sciuto: *Nucl. Phys. B* **410** (1993) 577
- [17] M. Rausch de Traubenberg, P. Simon: *Preprint* (hep-th/9806188)
- [18] L. P. Colatto, J. L. Matheus-Valle: *Preprint* (hep-th/9504101)
- [19] M. Daoud, Y. Hassouni: *Prog. Theor. Phys.* **96** (1997) 1033
- [20] M. Daoud, J. Douari, Y. Hassouni: *submitted to J. Math. Phys., Preprint* (IC/98/132)
- [21] M. Daoud, Y. Hassouni: *Mod. Phys. Lett. A* **96** (1997) 1033
- [22] M. Daoud, J. Douari, Y. Hassouni: *Int. J. Theor. Phys.* **36** (1997) 3071